

The genetics of early telencephalon patterning: some assembly required

Jean M. Hébert* and Gord Fishell†

Abstract | The immense range of human behaviours is rooted in the complex neural networks of the cerebrum. The creation of these networks depends on the precise integration of specific neuronal subtypes that are born in different regions of the telencephalon. Here, using the mouse as a model system, we review how these proliferative zones are established. Moreover, we discuss how these regions can be traced back in development to the function of a few key genes, including those that encode fibroblast growth factors (FGFs), sonic hedgehog (SHH), bone morphogenetic proteins (BMPs), forkhead box G1 (FOXP1), paired box 6 (PAX6) and LIM homeobox protein 2 (LHX2), that pattern the early telencephalon.

Neuroepithelium

The embryonic ectoderm that develops into the nervous system.

Neural plate

The neural epithelial cells that form in the early embryo after neuronal induction and give rise to the nervous system.

Anlagen

An embryonic precursor of a more mature tissue.

Morphogen

A secreted factor that can induce two or more different cell fates in a concentration-dependent manner by forming a gradient.

*Departments of Neuroscience and Molecular Genetics, Albert Einstein College of Medicine, 1410 Pelham Parkway South, Bronx, New York 10461, USA.
†Smilow Neuroscience Program and Department of Cell Biology, Smilow Research Building, New York University, 522 First Avenue, New York, New York 10016, USA.
e-mails:

jhebert@aecom.yu.edu;
fishell@saturn.med.nyu.edu

doi:10.1038/nrn2463

Published online 6 August 2008

Despite the complexity of the adult cerebrum, the telencephalon starts off as a simple neuroepithelium at the anterior end of the neural plate. The elaborate process by which this primordial sheet of cells progresses into a mature brain can be divided into several discrete phases and has been best studied in the mouse. The initial phases involve assigning anterior–posterior identity in the neuraxis, on the basis of antagonistic signalling between the organizer and the anterior visceral endoderm^{1–3}. Once the telencephalic anlagen is established, it is subdivided into different territories through the actions of morphogens that induce specific transcription factors. Following these early patterning events, the embryonic telencephalon gives rise to a dorsally positioned cortical ventricular zone and a series of ventral eminences that are positioned along the rostral–caudal axis. Each of these progenitor domains produces specific types of neurons that ultimately coalesce in the mature telencephalon and form connections that are then refined and modulated with experience. The neocortex is the major component of the dorsal telencephalon. It is largely comprised of excitatory glutamatergic principal cells (pyramidal neurons) and GABA (γ -aminobutyric acid)-ergic inhibitory interneurons. Pyramidal neurons are generated in the cortical ventricular proliferative zone. By contrast, interneurons originate in distinct ventral eminences and subsequently migrate dorsally into the developing cortex.

Here we focus exclusively on the steps in telencephalic development during which the telencephalic neuroepithelium is patterned into molecularly distinct proliferative zones that generate the neuronal diversity of the adult cerebrum. Specifically, we restrict

our attention to the events that take place between embryonic day (E) 8.5 and E11 in the mouse, after the telencephalic primordium has been specified. We outline the events in this period that are critical for later telencephalic patterning and neurogenesis. Numerous reviews have focused on the roles of specific signalling pathways and associated transcription factors, such as sonic hedgehog (SHH), fibroblast growth factors (FGFs) and members of the SIX3 and NKX families, in neural development^{4–7}. Moreover, anterior–posterior specification of the neuraxis, including the forebrain, has recently been considered^{3,8}. However, to date there have been few attempts to provide a description of the critical genetic interactions that link the extrinsic and intrinsic events by which the cerebrum is patterned. Here we examine recent insights into the genetic processes by which this is achieved.

Although some understanding of these events has been obtained from the study of lower vertebrates or invertebrates, we focus on findings in the mouse because the events that are unique to higher mammals are likely to be the most relevant for human telencephalic development. We start by discussing how the orchestrated interplay between extrinsic and intrinsic signals, including interactions between SHH and GLI3, GLI3 and forkhead box G1 (FOXP1), FOXP1 and FGFs, and FGFs and SHH (even in biology, what goes around comes around), results in the emergence of defined dorsal and ventral telencephalic territories. We then go on to examine how the major divisions are further refined, with a focus on the pallial–subpallial boundary and the dorsal subdivisions. We suggest that taken together these observations provide the basis for

Pallium

The roof of the telencephalon (but not synonymous with the cortex). It contains both cortical structures (for example, the hippocampus and the neocortex) and deep-lying nuclear structures (for example, the claustrum and parts of the amygdala).

Cortical hem

A transient structure located in the dorso-medial area of the embryonic telencephalon, between the hippocampal anlagen and the choroid plexus. The cortical hem acts as a hippocampal organizer.

a coherent model of telencephalic patterning. Our aim is to provide clarity in what is a complex field: readers should be aware that in many cases alternative interpretations certainly exist.

Early divisions in the telencephalon

Early in its development, the anterior neural plate gives rise to the prosencephalon, which is subsequently subdivided into the telencephalon and the diencephalon. The mechanisms that underlie these early processes have been reviewed elsewhere^{2,3,8}, so in this Review we confine ourselves to stages after the telencephalic primordium has been specified. This occurs at approximately E8.5 in mice, when expression of the forkhead transcription factor gene *Foxg1* is initiated in this region^{9,10}. At this stage the telencephalon is still a neuroepithelium of single-cell thickness. Immediately following *Foxg1* expression the telencephalon becomes subdivided into several distinct territories, which are first indicated by the expression of specific molecular markers and which shortly afterwards can be distinguished on the basis of regional differences in their levels of proliferation.

Genetic studies using model organisms such as the mouse and the zebrafish have identified the key transcription factors that regulate embryonic telencephalic patterning. These transcription factors define four broad areas that generate different cell types and develop into functionally distinct adult structures. The embryonic dorsal telencephalon generates primarily glutamatergic neurons and can be divided into two main regions: an anterior and lateral region that gives rise to the neocortex, and a posterior and medial area that develops into the hippocampus, the cortical hem

and the choroid plexus. The ventral telencephalon can also be divided into two main regions: a medial domain that is known as the medial ganglionic eminence (MGE), and two posterior and lateral regions that are designated the lateral ganglionic eminence (LGE) and the caudal ganglionic eminence (CGE). Each of these regions contributes neurons to discrete populations in the basal ganglia and to associated limbic structures, including the amygdala and the nucleus accumbens. In addition, the MGE produces *somatostatin* and *parvalbumin* subclasses of GABAergic interneurons (as well as a population of neuropeptide Y (NPY)-expressing interneurons) that ultimately reside in the basal ganglia and the cortex, whereas the CGE primarily produces *calretinin*- and/or vasoactive intestinal peptide (VIP)-expressing interneurons. By contrast, the LGE produces a substantial population of olfactory bulb interneurons, as well as inhibitory projection neurons that populate several ventral regions including the striatum and limbic areas^{11–13}.

The nascent anterior neural plate expresses several transcription factor genes in a region that includes the cells that are destined to form the telencephalon¹⁴. Among these genes are *Gli3*, *Foxg1* and paired box 6 (*Pax6*). As discussed below, each of these genes has a role in dividing the telencephalon into its dorsal and ventral sections.

SHH and GLI3 specify dorsal and ventral domains

The initial subdivision that defines what will later become the dorsal and the ventral telencephalon is regulated at least in part by the dorsalizing effects of *Gli3* expression and the ventralizing influence of *Shh* expression. *Gli3* encodes a zinc-finger transcription factor that was first identified as the gene that is disrupted in the classical mouse mutation ‘extra toes’ (*Xt*)¹⁵. *Gli3* is initially expressed broadly throughout the telencephalic anlagen and then is progressively downregulated in the ventral portion^{16,17} (FIG. 1). In the absence of *Gli3*, the development of the dorsal telencephalon is disrupted: the choroid plexus, the cortical hem and the hippocampus fail to form and the development of the neocortex is progressively compromised^{18–21}.

The *Shh* gene encodes a secreted signalling protein that is related to the *Drosophila* protein hedgehog. *Shh* is expressed in the midline of the nascent neural plate and continues to be expressed along the ventral midline of the CNS throughout development²² (FIG. 1). In *Shh*^{-/-} mouse embryos, the telencephalon is reduced in size and ventral cell types are lost^{17,23–25} (BOX 1). In double-mutant mice that lack both *Shh* and *Gli3*, however, ventral patterning is largely rescued^{16,26,27}. Therefore, SHH restricts the dorsalizing function of GLI3 and controls the positioning of the dorsoventral boundary. Hence, SHH promotes ventral identity by preventing dorsalization of the telencephalon, rather than by directly promoting ventral cell character. The rescue of ventral development in the *Shh* mutant through compound removal of *Gli3* indicates that other genes, acting independently or downstream of *Shh*, function positively to generate ventral cell types²⁶.




Figure 1 | Key extrinsic and intrinsic cellular factors establish the dorsal and ventral subdivisions of the telencephalon. Schematics of the anterior neural plate in a mouse embryo (dorsal view, five somites stage, anterior is up). **a** | The region that will become the telencephalon is defined by expression of forkhead box G1 (*Foxg1*) (shown in blue). **b** | FOXG1 and sonic hedgehog (SHH; green) promote fibroblast growth factor (FGF; purple) expression in the anterior neural ridge. This patterns the nascent telencephalon (indicated by the curved arrow). SHH promotes FGF expression indirectly by inhibiting the repressor activity of GLI3 (expression of GLI3 is shown in red). Consequently, SHH promotes the formation of a ventral telencephalic subdivision by inhibiting the dorsalizing effects of GLI3.

Box 1 | **Sonic hedgehog's role as a morphogen**

Morphogens are molecules that bestow cell identity in a concentration-dependent manner⁸¹. As such, they are known to be essential for CNS patterning. Of the extrinsic signals that have been implicated as morphogens, the regulation of the activity of sonic hedgehog (SHH) is the best understood. SHH signalling is modulated by numerous factors, including GLI effectors, which have been shown to be involved in establishing ventral identity throughout the developing CNS^{16,26,82}. GLI effectors come in two forms: constitutively cleaved forms that act as transcriptional repressors in the absence of SHH signalling, and uncleaved forms that function as transcriptional activators in the presence of SHH ligands. However, the precise utilization of different components of the SHH signalling pathway is level-specific. For instance, whereas GLI activators are critical to ventral patterning in the spinal cord, GLI repressors are primarily required in the telencephalon^{26,82}. In the forebrain, SHH signalling also requires other factors for ventral development, such as smoothed (SMO; the obligate downstream effector of SHH), low-density lipoprotein receptor-related protein 2 (LRP2, also known as megalin) and the multifunctional transmembrane protein CDO^{83–86}.

Gli3 and Foxg1 maintain the telencephalon

FOXG1, through its ability to induce expression of the extracellular signal FGF8 (see below), is probably the main source of positive regulation of ventral telencephalic identity. *Foxg1* is expressed in the early anterior plate cells that are destined to form the telencephalon^{10,28,29} (FIG. 1). Its expression is independent of that of *Shh*: although *Foxg1* expression is impaired in *Shh*^{-/-} mice, experiments using *Shh*^{-/-};*Gli3*^{-/-} double-mutant mice revealed that this impairment occurs as a result of increased GLI3 repressor function²⁷. Disruption of *Foxg1* expression results in a loss of ventral cell types^{30–32}. However, unlike in the *Shh*^{-/-} phenotype, ventral cells are not rescued in the *Foxg1*^{-/-} mutants by removal of *Gli3* (REF. 33). In fact, the telencephalon is completely lost in the *Foxg1*^{-/-};*Gli3*^{-/-} double mutant, indicating that *Gli3* and *Foxg1* are essential for generating and maintaining the dorsal and ventral subdivisions of the telencephalon, respectively (FIG. 2).

FOXG1 and FGFs act cooperatively

In generating the ventral telencephalon, FOXG1 acts in concert with FGF signalling. FOXG1 is required for *Fgf8* expression³² and, conversely, *Foxg1* expression might itself be regulated by FGF signalling. FGF8-soaked beads can ectopically induce expression of *Foxg1* (REF. 10), and *Foxg1* expression is reduced in *Fgf8*^{-/-} mutants³⁴, leading to the hypothesis that FGFs and FOXG1 form a positive-feedback loop. Whether FGF signalling is essential for *Foxg1* expression remains unclear owing to potential functional compensation by other members of the FGF family in the FGF signalling mutants that have been examined to date.

FGF signalling, like SHH signalling, is essential for the generation of ventral cell types in the telencephalon^{34–38}. Moreover, there are data that indicate that FGFs act in a dose-dependent manner to pattern the ventral telencephalon. Disruption of the gene that encodes FGF receptor 1 (*Fgfr1*) leads to a loss of expression of LIM homeobox protein 6 (LHX6) and LHX7, two LIM-domain transcription factors that are expressed in the MGE and are necessary for the differentiation of MGE-derived interneurons^{38–40}. Disruption of both *Fgfr1* and *Fgfr2* results in an even more severe loss of ventral

precursor cells³⁸. In these mutants, both the homeobox gene *Nkx2.1*, which is normally expressed in the MGE and is necessary for its development⁴¹, and *Gsh2*, which is necessary for LGE development^{42,43}, are no longer expressed. Similarly, the phenotypes of severely hypomorphic and null alleles of *Fgf8* reveal a progressive loss of ventral markers (including *Nkx2.1* expression) and associated structures with diminishing levels of *Fgf8* expression³⁴.

FGFs downstream of Shh specify ventral identity

Several lines of evidence indicate that FGFs operate downstream of *Shh* to generate ventral cell types. SHH maintains the expression of *Fgf3*, *Fgf8*, *Fgf15*, *Fgf17* and *Fgf18* in the anterior medial telencephalon, and FGF receptors are required for the ventralizing effect of SHH^{16,25,27,38}. Moreover, FGF8-soaked beads can induce ventral gene expression in cultured dorsal telencephalic explants when SHH signalling is inhibited²¹.

Why are ventral cells still generated in the *Shh*^{-/-};*Gli3*^{-/-} double mutant if SHH is required to maintain the expression of FGF genes? The answer again is that expression of these FGF genes is only indirectly promoted by SHH, through its ability to negatively regulate GLI3's repressor function (FIG. 1). In the *Shh*^{-/-} mutant there is no FGF expression, and ventral cell types are lost because of the unchecked repressive action of GLI3. In the *Shh*^{-/-};*Gli3*^{-/-} double mutant, FGF expression is no longer attenuated by GLI3, and so ventral development is restored. FGF expression is similarly expanded in both the *Gli3*^{-/-} and the *Shh*^{-/-};*Gli3*^{-/-} mutants^{16,19,21,27}, confirming that SHH promotes FGF expression indirectly by attenuating GLI3's repressor function. Loss of *Gli3* does not rescue the dorsalized phenotype that results from a loss of FGF receptors, consistent with the notion that GLI3 acts upstream of FGF signalling³⁸ (FIG. 1).




Figure 2 | The dorsal and ventral telencephalic regions are subdivided into four major domains.

The dorsal and ventral subdivisions of the embryonic mouse telencephalon at embryonic day (E) 9.0 (a), and the four broad subdivisions at E10.0 (b). In both schematics, dorsal is up, ventral is down. The *Gli3*-expressing dorsal region at E9.0 is split, by E10.0, into a bone morphogenetic protein (BMP)- and Wingless/Int protein (WNT)-expressing medial region and a more lateral cortical region that expresses countergradients of empty spiracles homeobox 2 (*Emx2*) and paired box 6 (*Pax6*). The ventral region at E9.0 is split, by E10.0, into medial *Nkx2.1*-expressing domains and lateral *Gsh2*-expressing domains. At E10.0 the expression domain of *Gsh2* overlaps with that of *Nkx2.1*; this is not represented for the sake of illustrative simplicity. Similarly, sonic hedgehog (*Shh*), fibroblast growth factor (FGF) and forkhead box G1 (*Foxg1*) expression at E10.0 is omitted.

Mitogen

An agent that induces mitosis, usually resulting in cell division.

Because SHH specifies fates in the ventral CNS in a dose-dependent manner, it meets the strict definition of a morphogen (BOX 1). However, in the telencephalon at least, it does so indirectly through its regulation of FGFs. Similarly, it seems that much of SHH's mitogenic effects in the telencephalon are also FGF-mediated. This indicates that FGFs are the obligate effectors of SHH signalling and, by implication, that FGFs are both mitogens and morphogens in their own right.

FGF signalling acts as a telencephalic organizer

The dose-dependent patterning of the telencephalon by FGF signalling extends beyond the ventral regions into the dorsal regions. In embryos in which FGF signalling is only partially reduced by ectopic expression of a dominant-negative receptor, or in moderately hypomorphic *Fgf8* mutants, the transcription-factor gradients in the cortical neuroepithelium shift anteriorly^{44,45}. In these mutants, the areal territories that are normally positioned in the more posterior regions of the cortex, such as the visual and sensory areas, are expanded at the

expense of anterior areal regions, such as the prelimbic cortex. Moreover, as the levels of FGF signalling are progressively reduced, loss of anterior cortical regions is accompanied by a diminution of the ventral telencephalon (FIG. 3). Although only the ventromedial cell types (those that arise from the MGE) are lost in *Fgfr1*^{-/-} mutants, in *Fgfr1*^{-/-};*Fgfr2*^{-/-} double mutants all ventral cells are lost^{38,46}. Similarly, in severely hypomorphic and null *Fgf8* mutants, not only are ventral precursors lost but the neocortex is also progressively diminished in size, with a preferential loss of anterior-lateral markers³⁴. Moreover, preliminary results indicate that when the expression of all three of the FGF receptor genes that are normally expressed in telencephalic precursors (*Fgfr1*, *Fgfr2* and *Fgfr3*) is lost, both the ventral telencephalon and the cerebral cortex are lost (J.H., unpublished observations). Taken together, these findings suggest that, as in other tissues (BOX 2), FGF signalling acts as an organizer for the telencephalon.

Pax6 establishes cortical regions

Although FOXG1 is absolutely required for the establishment of ventral identity in the telencephalon, it might also collaborate with FGFs in partitioning the dorsal telencephalon into the anterior-lateral neocortex and the posterior-medial territories. In *Foxg1*^{-/-} embryos, although the ventral telencephalon is most severely affected, the cortex is also reduced in size³⁰. This is due at least in part to the premature differentiation of precursor cells into layer-1 Cajal-Retzius neurons⁴⁷. However, it is also partially due to the loss of anterior lateral precursor cells, as the remaining cortical field seems to be caudalized in *Foxg1*^{-/-} mutants^{33,48}. Hence, both FOXG1 and FGFs are required for normal cortical development, despite their ventralizing functions.

What enables FOXG1 and FGFs to specify ventral telencephalon in one context and anterior-lateral cortex in another? The answer, at least in part, involves PAX6, a paired-box transcription factor that is essential for setting up the sharp border that distinguishes the ventral telencephalon from the dorsal telencephalon. The embryonic patterning role of PAX6 was initially identified following genetic mapping of the classical 'small eye' (*sey*) mouse mutant⁴⁹. At the neural plate stage *Pax6* is expressed throughout the telencephalic anlagen¹⁴. At the neural tube stage its expression is downregulated in the region that will become the ventral telencephalon, concomitant with the upregulation of *Nkx2.1* in this region¹⁷. As a result, the boundary between *Pax6* and *Nkx2.1* expression initially demarcates the dorsoventral border. Slightly later in development (at E9.5 in mice), the *Pax6*- and *Nkx2.1*-expressing regions become separated by a domain of *Gsh2* expression¹⁷ (FIG. 2). As a result, the pallial-subpallial boundary becomes defined by the intersection of *Pax6* and *Gsh2* expression (the expression domains overlap slightly).

Structurally, the pallial-subpallial boundary occurs at the interface between the ventral cortical primordium and the most dorsal portion of the subpallium, designated the dorsal LGE. This is a remarkable boundary: at this border, the ordered laminar arrangement of the




Figure 3 | FGF signalling as an organizer for the telencephalon. Schematics of coronal and sagittal sections in the telencephalon between embryonic day (E) 12.5 and E15.5 in various mutant mice. The colours represent areas that contain cells with different telencephalic identities. Progressively deleting more fibroblast growth factor receptor (FGFR) genes specifically in the anterior neural plate leads to diminished FGF signalling and gradually more severe truncations of telencephalic regions. Anterior ventral regions are lost first, followed by posterior dorsal regions^{34,38,46}.

Box 2 | FGFs mediate organizer activity in multiple developmental contexts

The experiments conducted by Spemann and Mangold in 1924 defined an organizer as a group of cells that can induce neighbouring cells to change their fate and reorganize into a new, normally patterned tissue⁸⁷. Fibroblast growth factors (FGFs), and in particular FGF8, are expressed in several organizers in the developing embryo, including the apical ectodermal ridge (at the tips of the limb buds) and the isthmus at the midbrain–hindbrain boundary⁸⁸. In these locations FGF8 itself is likely to mediate organizer activity, as it is both necessary and sufficient (when it is ectopically expressed) to induce the formation of a normally structured limb and midbrain^{89–95}. FGFs are also expressed in the anterior neural ridge (FIG. 1), a group of cells that forms the border between the most anterior part of the ectoderm and the neuroectoderm and that patterns the telencephalon^{10,88,96}. The loss of parts of the telencephalon that is observed in mutants that lack combinations of the three FGF receptors in the anterior neural plate suggests that FGFs that emanate from the anterior neural ridge might mediate organizer activity for the telencephalon³⁸. These results, together with some preliminary findings in mutants lacking *Fgfr1*, *Fgfr2* and *Fgfr3* (J.H., unpublished observations), indicate that, unlike in the midbrain–hindbrain boundary and the limb, loss of FGF8 can be compensated for by other FGF ligands in organizing the telencephalon^{34,38}. FGFs are also likely to be sufficient to induce the telencephalon in the context of the early anterior neural plate, as FGF8 can induce forkhead box G1 (*Foxg1*) expression in the early anterior neural plate¹⁰ and can cause mirror-image duplications (a hallmark of organizer activity) of somatosensory whisker barrels later in the development of the dorsal telencephalon⁴⁴. Hence, the combined effect of several FGFs that emanate from the anterior ridge seems to mediate organizer activity to generate and pattern all but the posterior medial telencephalon (FIG. 3).

neurons in the cortex shifts to the apparently chaotic nuclear structure of the subpallium. *Pax6* and *Gsh2* help to establish this boundary and show a mirrored pattern of expression, with *Pax6* being expressed in a dorsal (low) to ventral (high) gradient and *Gsh2* being expressed in a ventral (low) to dorsal (high) gradient. Loss-of-function analysis of these two genes demonstrates that their mutual antagonism is required for the positioning of the boundary. In *Pax6*^{-/-} embryos the most ventral area of the cortical primordium becomes the dorsal LGE, whereas in *Gsh2*^{-/-} embryos the reverse occurs and the dorsal LGE is transformed into the ventral cortex^{50–53}.

In a phenotype that is reminiscent of the broader rescue of dorsoventral patterning that is seen in *Gli3*^{-/-};*Shh*^{-/-} double mutants, patterning of the pallial–subpallial boundary is partially restored in *Pax6*^{-/-};*Gsh2*^{-/-} double mutant embryos. This suggests that these genes form only part of a broader genetic network that maintains the positioning of this boundary⁵². Consistent with this notion, loss of *Pax6* function, like loss of *Gli3*, partially compensates for the loss of *Shh*, suggesting that GLI3 and PAX6 interact to promote dorsal fates⁵⁴. In fact, although GLI3 is not necessary for initial *Pax6* expression, it is partially required to maintain *Pax6* expression^{16,19,21}. Determining whether loss of *Pax6* can suppress the dorsalization of the telencephalon that is observed in FGF-gene mutants and *Foxg1* mutants should help us to elucidate *Pax6*'s position in the dorsal–ventral regulatory cascade. Moreover, even though SHH and FGFs seem to be the key ventralizing signals, it is evident that other factors, such as TLX and retinoic acid, probably participate in setting up the boundary between the cortex and the ventral telencephalon^{55,56}.

Some of the cooperative dorsalizing actions of PAX6 and GLI3 seem to be mediated through empty spiracles homeobox 2 (*EMX2*), a transcription factor that is related to empty spiracles protein. With regards to cortical patterning, *EMX2* seems to lie genetically downstream of GLI3, as indicated by the loss of *Emx2* expression in *Gli3* mutants⁵⁷. Furthermore, *Pax6*^{-/-};*Gli3*^{-/-} double mutants resemble *Pax6*^{-/-};*Emx2*^{-/-} double mutants, in that both compound mutants exhibit an expansion of the cortical

hem and the subpallium at the expense of the isocortex^{54,58}. One copy of any of these genes in either compound mutant is sufficient to maintain cortical identity.

In addition to acting cooperatively to specify telencephalic precursor cells as cortical, *EMX2* and *PAX6* act individually and antagonistically to further pattern the cortical neuroepithelium^{59–63}. These proteins, among numerous other factors including the nuclear receptors coup-TF1 and TLX, the zinc-finger transcription factor SP8 and the basic helix–loop–helix proteins neurogenin 1 and 2, are also likely to be important in patterning cortical precursors^{35,64–70}. The fact that certain factors are required for both cortical specification and progenitor proliferation underscores the need for developmental coordination of areal and laminar patterning in the assembly of cortical architecture.


Partitioning the dorsal telencephalon

The dorsal telencephalon is split into two broad domains: the cerebral cortex, which gives rise mainly to the neocortex and the hippocampus; and the dorsal midline, which gives rise to the choroid plexus and the cortical hem. *LHX2*, a LIM homeodomain transcription factor that is expressed in the cortical primordium, is crucial in specifying cells as cortical rather than dorsal midline. In *Lhx2*^{-/-} embryos, the entire area that normally becomes the cortex is lost at the expense of an expanded choroid plexus and cortical hem⁷¹. In fact, experiments that used chimaeras and mosaics demonstrated that *Lhx2* acts as a classic selector gene to specify a cortical fate and inhibit a hem or anti-hem fate⁷² (BOX 3). *FOXG1* also has a role in confining the development of the dorsal midline, as in *Foxg1*^{-/-} mutants dorso-medial markers (such as bone morphogenetic protein 4 (*Bmp4*) expression) are expanded well beyond the length of their normal domain^{31,32}.

What promotes the development of the dorsal midline itself? The dorsal midline is marked by the expression of two families of secreted signalling molecules: BMPs and Wingless/Int proteins (WNTs). BMP signalling can induce dorsal midline features both *in vitro* and *in vivo*. BMP-soaked beads placed on explants of cortical

Box 3 | Signals from the midline subdivide the dorsal telencephalon

Bone morphogenetic proteins (BMPs), which are expressed in midline cells, are required for the formation of the choroid plaque (the precursor to the choroid plexus) and the cortical hem^{74,75}. In turn, the cortical hem acts as an organizer for the hippocampus⁷². Wingless/INT proteins (WNTs) are likely to be important components of this organizer activity, as without secretion of WNT3a from the hem the hippocampus fails to develop⁷⁷. Intracellular effectors of BMP and WNT signalling can bind to enhancer elements found upstream of empty spiracles homologue 2 (*Emx2*) and regulate its expression. WNT3a might therefore act together with BMPs to promote hippocampal formation by directly regulating the expression of *EMX* genes^{57,76}. The neocortex, however, requires both *EMX*-gene expression and *Pax6* expression to form⁵⁸. In addition, LIM homeobox protein 2 (*Lhx2*) acts as a selector gene that cell-autonomously specifies cortical fates, both hippocampal and neocortical⁷². *LHX2* also suppresses the infringing of flanking cell fates, by repressing expansion of the cortical hem and antihem, a band of cells at the lateral extreme of the pallium⁹⁷. Note that *Pax6*, which is only partially regulated by *LHX2*, is required for formation of the antihem. At present the signals that regulate the expression of *Lhx2* and *Pax6* remain unknown. One candidate for a source of such signals is the antihem, which might be involved in patterning the dorsal telencephalon. These signalling pathways are illustrated in the figure.



neuroepithelium induce midline features, such as apoptosis and *Msx1* expression, and repress lateral features, such as proliferation and *Foxg1* expression⁷³. In addition, transgenic expression of an activated BMP receptor in the cortical neuroepithelium leads to an expansion of midline cell types at the expense of cortical ones⁷⁴. Moreover, BMP signalling is required for the formation of the dorsal midline, as in mice that lack both BMP receptor 1a and BMP receptor 1b (*Bmpr1a*^{-/-}; *Bmpr1b*^{-/-} double mutants), the choroid plexus and cortical hem fail to form⁷⁵.




Figure 4 | Key genetic pathways that interact to form and pattern the early telencephalon. Factors that act early to establish broad telencephalic regions are shown in blue. Sonic hedgehog (SHH) ventralizes the telencephalon by antagonizing the dorsalizing effect of GLI3. By repressing *Gli3*, SHH, together with forkhead box G1 (FOXG1), activates fibroblast growth factor (FGF) expression. FGF might feedback and promote *Foxg1* expression (dotted arrow). FOXG1 and FGF signalling are necessary for forming all regions of the telencephalon (shown in green), except for the dorsomedial region (shown in orange). Downstream transcription factors, such as GSH2 and NKX2.1, then form specific subdivisions. In the dorsal telencephalon, GLI3's promotion of the expression of bone morphogenetic proteins (BMPs) and Wingless/Int proteins (WNTs) is required for *EMX*-gene expression. The products of the *EMX* genes, along with PAX6 and LHX2, further subdivide the dorsal telencephalon. LGE, lateral ganglionic eminence; MGE, medial ganglionic eminence.

In *Gli3*^{-/-} embryos the dorsal midline also fails to form, presumably because the expression of BMP genes is lost^{18–21}. Immediately adjacent to the dorsal midline is the hippocampal primordium, the development of which requires *EMX1* and *EMX2* (REF. 76). *Emx1* and *Emx2* expression throughout the cortical primordium depends on *GLI3*^{19,20}. This is likely to be an indirect dependency that requires BMPs and WNTs that emanate from the cortical hem, which acts as a hippocampal organizer^{57,72,77,78}. Conversely, *EMX2* might feedback and promote WNT-gene expression^{79,80}. Hence, as with the other broad domains of the developing cerebral hemispheres, key players have been identified that regulate the development of dorso-medial structures.

Conclusion

Much progress has been made in the past decade in understanding the genetic pathways that are involved in patterning the early telencephalon. A small number of intrinsic and extrinsic cellular factors have been identified that interact to set up discrete telencephalic domains (FIG. 4). To fully understand the mechanisms behind telencephalic development, it will be necessary to further elucidate how the known patterning factors regulate each other's function, what additional factors are involved, how multiple signals are integrated over time in precursor cells, and how this allows them to progressively restrict their identity. It is likely that additional roles for factors such as WNTs and BMPs will be revealed. Collectively, these developments will lead to an understanding of how the various factors impart individual neurons with appropriate regional and cell-type identities in closely adjacent telencephalic territories. Appreciation of the mechanisms that underlie the assembly of the telencephalon will also allow us to gain further insights into the molecular underpinnings of developmental brain disorders. A deeper understanding of how the fate of neural precursor cells is regulated *in vivo* will undoubtedly provide new diagnostic and prognostic tools and therapeutic targets for treating these conditions.

1. Beddington, R. S. & Robertson, E. J. Anterior patterning in mouse. *Trends Genet.* **14**, 277–284 (1998).
2. Wilson, S. W. & Rubenstein, J. L. Induction and dorsoventral patterning of the telencephalon. *Neuron* **28**, 641–651 (2000).
3. Rallu, M., Corbin, J. G. & Fishell, G. Parsing the prosencephalon. *Nature Rev. Neurosci.* **3**, 5031–5040 (2002).
4. Ingham, P. & McMahon, A. P. Hedgehog signaling in animal development: paradigms and principles. *Genes Dev.* **15**, 3059–3087 (2001).
5. Kobayashi, D. *et al.* Early subdivisions in the neural plate define distinct competence for inductive signals. *Development* **129**, 83–93 (2002).
6. Fuccillo, M., Joyner, A. L. & Fishell, G. Morphogen to mitogen: the multiple roles of hedgehog signalling in vertebrate neural development. *Nature Rev. Neurosci.* **7**, 772–783 (2006).
7. Mason, I. Initiation to end point: the multiple roles of fibroblast growth factors in neural development. *Nature Rev. Neurosci.* **8**, 583–596 (2007).
8. Wilson, S. W. & Houart, C. Early steps in the development of the forebrain. *Dev. Cell* **6**, 167–181 (2004).
9. Tao, W. & Lai, E. Telencephalon-restricted expression of BF-1, a new member of the HNF-3/*fork head* gene family, in the developing rat brain. *Neuron* **8**, 957–966 (1992).
This paper provided the first description of *Foxg1*^{-/-} mice. It supplied strong evidence that *Foxg1* is centrally involved in the specification of the telencephalon.
10. Shimamura, K. & Rubenstein, J. L. Inductive interactions direct early regionalization of the forebrain. *Development* **124**, 2709–2718 (1997).
This beautiful paper provided the first evidence that FGF8 can induce FOXG1 signalling and hence the specification of the telencephalon.
11. Wichterle, H., Turnbull, D. H., Nery, S., Fishell, G. & Alvarez-Buylla, A. *In utero* fate mapping reveals distinct migratory pathways and fates of neurons born in the mammalian basal forebrain. *Development* **128**, 3759–3771 (2001).
12. Nery, S., Fishell, G. & Corbin, J. G. The caudal ganglionic eminence is a source of distinct cortical and subcortical cell populations. *Nature Neurosci.* **5**, 1279–1287 (2002).
13. Wonders, C. P. & Anderson, S. A. The origin and specification of cortical interneurons. *Nature Rev. Neurosci.* **7**, 687–696 (2006).
14. Inoue, T., Nakamura, S. & Osumi, N. Fate mapping of the mouse prosencephalic neural plate. *Dev. Biol.* **219**, 373–383 (2000).
15. Hui, C.-C. & Joyner, A. L. A mouse model of Greig cephalo-polysyndactyly syndrome: the *extra-toes*['] mutation contains an intragenic deletion of the *Gli3* gene. *Nature Genet.* **3**, 241–246 (1993).
16. Aoto, K., Nishimura, T., Eto, K. & Motoyama, J. Mouse GLI3 regulates *Fgf8* expression and apoptosis in the developing neural tube, face, and limb bud. *Dev. Biol.* **251**, 320–332 (2002).
The paper, which was published concurrently with reference 26, demonstrated that SHH signalling in the telencephalon represses the expansion of dorsal signals rather than directly specifies ventral identity (a role that seems to be primarily subserved by FGF signals).
17. Corbin, J. G., Rutlin, M., Gaiano, N. & Fishell, G. Combinatorial function of the homeodomain proteins Nkx2.1 and Gsh2 in ventral telencephalic patterning. *Development* **130**, 4895–4906 (2003).
18. Grove, E. A., Tole, S., Limon, J., Yip, L. & Ragsdale, C. W. The hem of the embryonic cerebral cortex is defined by the expression of multiple *Wnt* genes and is compromised in *Gli3*-deficient mice. *Development* **125**, 2315–2325 (1998).
This was the first of a series of papers by this group which were seminal in elucidating the patterning of the cortical hem. Indeed, this paper was the first to introduce the concept of the cortical hem as a hippocampal organizing centre.
19. Theil, T., Alvarez-Bolado, G., Walter, A. & Ruther, U. *Gli3* is required for *Emx* gene expression during dorsal telencephalon development. *Development* **126**, 3561–3571 (1999).
20. Tole, S., Ragsdale, C. W. & Grove, E. A. Dorsoventral patterning of the telencephalon is disrupted in the mouse mutant *extra-toes*. *Dev. Biol.* **217**, 254–265 (2000).
21. Kuschel, S., Ruther, U. & Theil, T. A disrupted balance between Bmp/Wnt and Fgf signaling underlies the ventralization of the *Gli3* mutant telencephalon. *Dev. Biol.* **260**, 484–495 (2003).
22. Echelard, Y. *et al.* Sonic hedgehog, a member of a family of putative signaling molecules, is implicated in the regulation of CNS polarity. *Cell* **75**, 1417–1430 (1993).
23. Ericson, J. *et al.* Sonic hedgehog induces the differentiation of ventral forebrain neurons: a common signal for ventral patterning within the neural tube. *Cell* **81**, 747–756 (1995).
This paper provided the first indication that SHH signalling is directly required in the patterning of the ventral forebrain, in a manner that is analogous to its action in the spinal cord.
24. Chiang, C. *et al.* Cyclopa and defective axial patterning in mice lacking Sonic hedgehog gene function. *Nature* **383**, 407–413 (1996).
This paper provided the first description of the *Shh*^{-/-} mutant and demonstrated that SHH is required for the establishment of ventral patterning throughout the nervous system, as previously predicted by the McMahon–Tabin and Jessell groups.
25. Ohkubo, Y., Chiang, C. & Rubenstein, J. L. Coordinate regulation and synergistic actions of BMP4, SHH and FGF8 in the rostral prosencephalon regulate morphogenesis of the telencephalic and optic vesicles. *Neuroscience* **111**, 1–17 (2002).
26. Rallu, M. *et al.* Dorsoventral patterning is established in the telencephalon of mutants lacking both *Gli3* and Hedgehog signaling. *Development* **129**, 4963–4974 (2002).
This paper showed that ventral patterning in the telencephalon can be established independently of SHH signalling. It demonstrated that the dorsalization that is seen in *Shh* mutants is largely a result of increased GLI3 activator function. See also reference 16.
27. Rash, B. G. & Grove, E. A. Patterning the dorsal telencephalon: a role for Sonic hedgehog? *J. Neurosci.* **27**, 11595–11603 (2007).
28. Shimamura, K., Hartigan, D. J., Martinez, S., Puelles, L. & Rubenstein, J. L. Longitudinal organization of the anterior neural plate and neural tube. *Development* **121**, 3923–3933 (1995).
29. Hébert, J. M. & McConnell, S. K. Targeting of *cre* to the *Foxg1* (*BF-1*) locus mediates *loxP* recombination in the telencephalon and other developing head structures. *Dev. Biol.* **222**, 296–306 (2000).
30. Xuan, S. *et al.* Winged helix transcription factor BF-1 is essential for the development of the cerebral hemispheres. *Neuron* **14**, 1141–1152 (1995).
31. Dou, C.-L., Li, S. & Lai, E. Dual role of brain factor-1 in regulating growth and patterning of the cerebral hemispheres. *Cereb. Cortex* **9**, 543–550 (1999).
32. Martynoga, B., Morrison, H., Price, D. J. & Mason, J. O. *Foxg1* is required for specification of ventral telencephalic and region-specific regulation of dorsal telencephalic precursor proliferation and apoptosis. *Dev. Biol.* **283**, 113–127 (2005).
33. Hanashima, C., Fernandes, M., Hébert, J. M. & Fishell, G. The role of *Foxg1* and dorsal midline signaling in the generation of Cajal–Retzius subtypes. *J. Neurosci.* **27**, 11103–11111 (2007).
34. Storm, E. *et al.* Dosage dependent functions of *Fgf8* in regulating telencephalic patterning centers. *Development* **133**, 1831–1844 (2006).
35. Shanmugalingam, S. *et al.* *Ace/Fgf8* is required for forebrain commissure formation and patterning of the telencephalon. *Development* **127**, 2549–2561 (2000).
36. Shinya, M., Koshida, S., Sawada, A., Kuroiwa, A. & Takeda, H. Fgf signalling through MAPK cascade is required for development of the subpallial telencephalon in zebrafish embryos. *Development* **128**, 4153–4164 (2001).
37. Walshe, J. & Mason, I. Unique and combinatorial functions of *Fgf3* and *Fgf8* during zebrafish forebrain development. *Development* **130**, 4337–4349 (2003).
38. Gutin, G. *et al.* FGF acts independently of SHH to generate ventral telencephalic cells. *Development* **133**, 2937–2946 (2006).
This paper provided the first definitive evidence that FGF signalling is the primary positive effector that establishes ventral identity in the telencephalon. The findings in this paper were complemented by those in reference 34, which demonstrated that FGF8 is central to this process.
39. Fragkouli, A. *et al.* Loss of forebrain cholinergic neurons and impairment in spatial learning and memory in LHX7-deficient mice. *Eur. J. Neurosci.* **21**, 2923–2938, (2005).
40. Liodis, P. *et al.* *Lhx6* activity is required for the normal migration and specification of cortical interneuron subtypes. *J. Neurosci.* **27**, 3078–3089 (2007).
41. Sussel, L., Marin, O., Kimura, S. & Rubenstein, J. L. Loss of *Nkx2.1* homeobox gene function results in a ventral to dorsal molecular respecification within the basal telencephalon: evidence for a transformation of the pallidum into the striatum. *Development* **126**, 3359–3370 (1999).
42. Toresson, H. & Campbell, K. A role for *Gsh1* in the developing striatum and olfactory bulb of *Gsh2* mutant mice. *Development* **128**, 4769–4780 (2001).
43. Yun, K., Garel, S., Fischman, S. & Rubenstein, J. L. R. Patterning of the lateral ganglionic eminence by the *Gsh1* and *Gsh2* homeobox genes is required for histogenesis of the striatum and olfactory bulb and the growth of axons through the basal ganglia. *J. Comp. Neurol.* **461**, 151–165 (2003).
44. Fukuchi-Shimogori, T. & Grove, E. A. Neocortex patterning by the secreted signaling molecule FGF8. *Science* **294**, 1071–1074 (2001).
45. Garel, S., Huffman, K. J. & Rubenstein, J. L. Molecular regionalization of the neocortex is disrupted in *Fgf8* hypomorphic mutants. *Development* **130**, 1903–1914 (2003).
46. Hébert, J. M., Lin, M., Partanen, J., Rossant, J. & McConnell, S. K. FGF signaling through FGFR1 is required for olfactory bulb morphogenesis. *Development* **130**, 1101–1111 (2003).
47. Hanashima, C., Li, S. C., Shen, L., Lai, E. & Fishell, G. *Foxg1* suppresses early cortical cell fate. *Science* **303**, 56–59 (2004).
48. Muzio, L. & Mallamaci, A. J. *Foxg1* confines Cajal–Retzius neuronogenesis and hippocampal morphogenesis to the dorsomedial pallidum. *J. Neurosci.* **25**, 4435–4441 (2005).
49. Hill, R. E. *et al.* Mouse *Small eye* results from mutations in a paired-like homeobox-containing gene. *Nature* **354**, 522–525 (1991).
50. Corbin, J. G., Gaiano, N., Machold, R. P., Langston, A. & Fishell, G. The *Gsh2* homeodomain gene controls multiple aspects of telencephalic development. *Development* **127**, 5007–5020 (2000).
51. Stoykova, A., Treichel, D., Hallonet, M. & Gruss, P. *Pax6* modulates the dorsoventral patterning of the mammalian telencephalon. *J. Neurosci.* **20**, 8042–8050 (2000).
52. Toresson, H., Potter, S. & Campbell, K. Genetic control of dorsal-ventral identity in the telencephalon: opposing roles for *Pax6* and *Gsh2*. *Development* **127**, 4361–4371 (2000).
53. Yun, K., Potter, S. & Rubenstein, J. L. R. *Gsh2* and *Pax6* play complementary roles in dorsoventral patterning of the mammalian telencephalon. *Development* **128**, 193–205 (2001).
54. Fuccillo, M., Rutlin, M. & Fishell, G. Removal of *Pax6* partially rescues the loss of ventral structures in *Shh* null mice. *Cereb. Cortex* **16** (Suppl. 1) i96–i102 (2006).
55. Stenman, J. M., Wang, B. & Campbell, K. *Tlx* controls proliferation and patterning of lateral telencephalic progenitor domains. *J. Neurosci.* **23**, 10568–10576 (2003).
56. Marklund, M. *et al.* Retinoic acid signalling specifies intermediate character in the developing telencephalon. *Development* **131**, 4323–4332 (2004).
57. Theil, T., Aydin, S., Koch, S., Grotewold, L. & Ruther, U. Wnt and Bmp signalling cooperatively regulate graded *Emx2* expression in the dorsal telencephalon. *Development* **129**, 3045–3054 (2002).
This is a rare example of a study that examined a direct link between extracellular signals and the regulation of key transcription factors; the authors found that effectors of BMP and WNT signalling directly bind upstream cis elements at the *Emx2* locus to regulate its expression.
58. Muzio, L. *et al.* Conversion of cerebral cortex into basal ganglia in *Emx2*^{-/-} *Pax6*^{Shy/Sev} double-mutant mice. *Nature Neurosci.* **5**, 737–745 (2002).
59. Bishop, K. M., Goudreau, G. & O’Leary, D. D. M. Regulation of area identity in the mammalian neocortex by *Emx2* and *Pax6*. *Science* **288**, 344–349 (2000).
This paper provided the first attempt to explain how expression gradients of two transcription factors, EMX2 and PAX6, establish area identity (that is, functional territories) in the telencephalon. Coup-TF1 has also now been shown to contribute to this process (see reference 69). Recently this model has been codified into the ‘cooperative concentration’ model (see reference 63).

60. Mallamaci, A., Muzio, L., Chan, C.-H., Parnavelas, J. & Boncinelli, E. Area identity shifts in the early cerebral cortex of *Emx2*^{-/-} mutant mice. *Nature Neurosci.* **3**, 679–686 (2000).
This was the first paper to show that the loss of a specific transcription factor results in a shift in areal identity in the neocortex.
61. Bishop, K. M., Rubenstein, J. L. & O'Leary, D. D. Distinct actions of *Emx1*, *Emx2*, and *Pax6* in regulating the specification of areas in the developing neocortex. *J. Neurosci.* **22**, 7627–7638 (2002).
62. Muzio, L. *et al.* *Emx2* and *Pax6* control regionalization of the pre-neurogenic cortical primordium. *Cereb. Cortex* **12**, 129–139 (2002).
63. Hamasaki, T., Leingartner, A., Ringstedt, T. & O'Leary, D. D. *EMX2* regulates sizes and positioning of the primary sensory and motor areas in neocortex by direct specification of cortical progenitors. *Neuron* **43**, 359–372 (2004).
64. Zhou, C., Tsai, S. Y. & Tsai, M. J. *COUP-TFI*: an intrinsic factor for early regionalization of the neocortex. *Genes Dev.* **15**, 2054–2059 (2001).
65. Schuurmans, C. & Guillemot, F. Molecular mechanisms underlying cell fate specification in the developing telencephalon. *Curr. Opin. Neurobiol.* **12**, 26–34 (2002).
66. Stenman, J., Yu, R. T., Evans, R. M. & Campbell, K. *Tlx* and *Pax6* co-operate genetically to establish the pallio-subpallial boundary in the embryonic mouse telencephalon. *Development* **130**, 1113–1122 (2003).
67. Roy, K. *et al.* The *Tlx* gene regulates the timing of neurogenesis in the cortex. *J. Neurosci.* **24**, 8333–8345 (2004).
68. Schuurmans, C. *et al.* Sequential phases of cortical specification involve *Neurogenin*-dependent and -independent pathways. *EMBO J.* **23**, 2892–2902 (2004).
69. Armentano, M. *et al.* *COUP-TFI* regulates the balance of cortical patterning between frontal/motor and sensory areas. *Nature Neurosci.* **10**, 1277–1286 (2007).
70. Zembrzycki, A., Griesel, G., Stoykova, A. & Mansouri, A. Genetic interplay between the transcription factors *Sp8* and *Emx2* in the patterning of the forebrain. *Neural Develop.* **2**, 8 (2007).
71. Monuki, E. S., Porter, F. D. & Walsh, C. A. Patterning of the dorsal telencephalon and cerebral cortex by a roof plate-*Lhx2* pathway. *Neuron* **32**, 591–604 (2001).
72. Mangale, V. S. *et al.* *Lhx2* selector activity specifies cortical identity and suppresses hippocampal organizer fate. *Science* **319**, 304–309 (2008).
Although previous work by this group and others had demonstrated a requirement for LHX2 in the development of the neocortex, this paper definitively demonstrated a complete failure in neocortical specification in *Lhx2*^{-/-} mice. The finding suggests that LHX2 acts high in the hierarchy of neocortical induction, and provides an entry point for understanding neocortical specification.
73. Furuta, Y., Piston, D. W. & Hogan, B. L. M. Bone morphogenetic proteins (BMPs) as regulators of dorsal forebrain development. *Development* **124**, 2203–2212 (1997).
74. Panchision, D. M. *et al.* Sequential actions of BMP receptors control neural precursor cell production and fate. *Genes Dev.* **15**, 2094–2110 (2001).
75. Fernandes, M., Gutin, G., Alcorn, H., McConnell, S. K. & Hébert, J. M. Mutations in the BMP pathway in mice support the existence of two molecular classes of holoprosencephaly. *Development* **134**, 3789–3794 (2007).
76. Shinozaki, K., Yoshida, M., Nakamura, M., Aizawa, S. & Suda, Y. *Emx1* and *Emx2* cooperate in initial phase of archipallium development. *Mech. Dev.* **121**, 475–489 (2004).
77. Lee, S. M. K., Tole, S., Grove, E. & McMahon, A. P. A local *Wnt3a* signal is required for development of the mammalian hippocampus. *Development* **127**, 457–467 (2000).
78. Backman, M. *et al.* Effects of canonical Wnt signaling on dorso-ventral specification of the mouse telencephalon. *Dev. Biol.* **279**, 155–168 (2005).
79. Shimogori, T., Banuchi, V., Ng, H. Y., Strauss, J. B. & Grove, E. A. Embryonic signaling centers expressing BMP, WNT and FGF proteins interact to pattern the cerebral cortex. *Development* **131**, 5639–5647 (2004).
80. Muzio, L., Soria, J. M., Pannese, M., Piccolo, S. & Mallamaci, A. A mutually stimulating loop involving *Emx2* and canonical Wnt signalling specifically promotes expansion of occipital cortex and hippocampus. *Cereb. Cortex* **15**, 2021–2028 (2005).
81. Turing, A. M. The chemical basis of morphogenesis. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **237**, 37–72 (1952).
82. Bai, C. B., Stephen, D. & Joyner, A. L. All mouse ventral spinal cord patterning by hedgehog is *Gli3* dependent and involves an activator function of *Gli3*. *Dev. Cell* **6**, 103–115 (2004).
83. Fuccillo, M., Rallu, M., McMahon, A. P. & Fishell, G. Temporal requirement for hedgehog signaling in ventral telencephalic patterning. *Development* **131**, 5031–5040 (2004).
84. Spoelgen, R. *et al.* *LRP2/megalin* is required for patterning of the ventral telencephalon. *Development* **132**, 405–414 (2005).
85. Zhang, W., Kang, J. S., Cole, F., Yi, M.-J. & Krauss, R. S. *Cdo* functions at multiple points in the Sonic hedgehog pathway, and *Cdo*-deficient mice accurately model human holoprosencephaly. *Dev. Cell* **10**, 657–665 (2006).
86. Fernandes, M. & Hébert, J. M. The ups and downs of holoprosencephaly, dorsal versus ventral patterning forces. *Clin. Genet.* **73**, 413–423 (2008).
87. Spemann, H. & Mangold, H. Über induktion von Embryonalanlagen durch Implantation artfremder Organisatoren. *Wilhelm Roux. Arch. Entwicklungsmech. Organ.* **100**, 599–638 (1924).
88. Crossley, P. H. & Martin, G. R. The mouse *Fgf8* gene encodes a family of polypeptides and is expressed in regions that direct outgrowth and patterning in the developing embryo. *Development* **121**, 439–451 (1995).
89. Niswander, L., Tickle, C., Vogel, A., Booth, I. & Martin, G. R. *FGF-4* replaces the apical ectodermal ridge and directs outgrowth and patterning of the limb. *Cell* **75**, 579–587 (1993).
90. Lewandoski, M., Sun, X. & Martin, G. R. *Fgf8* signalling from the AER is essential for normal limb development. *Nature Genet.* **26**, 460–463 (2000).
91. Sun, X., Mariani, F. V. & Martin, G. R. Functions of FGF signalling from the apical ectodermal ridge in limb development. *Nature* **418**, 501–508 (2002).
92. Chi, C. L., Martinez, S., Wurst, W. & Martin, G. R. The isthmic organizer signal *FGF8* is required for cell survival in the prospective midbrain and cerebellum. *Development* **130**, 2633–2644 (2003).
93. Crossley, P. H., Martinez, S. & Martin, G. R. Midbrain development induced by *FGF8* in the chick embryo. *Nature* **380**, 66–68 (1996).
94. Crossley, P. H., Minowada, G., MacArthur, C. A. & Martin, G. R. Roles for *FGF8* in the induction, initiation, and maintenance of chick limb development. *Cell* **84**, 127–136 (1996).
95. Sato, T., Joyner, A. L. & Nakamura, H. How does *Fgf* signaling from the isthmic organizer induce midbrain and cerebellum development? *Dev. Growth Differ.* **46**, 487–494 (2004).
96. Houart, C., Westerfield, M. & Wilson, S. W. A small population of anterior cells patterns the forebrain during zebrafish gastrulation. *Nature* **391**, 788–792 (1998).
97. Assimakopoulos, S., Grove, E. A. & Ragsdale, C. W. Identification of a *Pax6*-dependent epidermal growth factor family signaling source at the lateral edge of the embryonic cerebral cortex. *J. Neurosci.* **23**, 6399–6403 (2003).

Acknowledgements

We would like to apologize to the authors of the many studies that have not been mentioned owing to lack of space. We would like to thank the many members of the Hébert and Fishell laboratories who read and offered suggestions on this Review, especially Drs Rossignol, Machold, Sousa, Demarco and Karayannis. In addition we would like to thank J. Dasen and B. Rudy for their comments on earlier versions of this manuscript. This research was supported by US National Institutes of Health grants to J.H. and G.F., as well as by funding from the Simons Foundation to G.F. and from the McDonnell and Sinsheimer Foundations to J.H.

DATABASES

Entrez Gene: <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene>
[calretinin](#) | [EMX2](#) | [FGF8](#) | [Fgfr1](#) | [FOXG1](#) | [GLI3](#) | [Gsh2](#) | [LHX6](#) | [LHX7](#) | [Nkx2.1](#) | [NPY](#) | [parvalbumin](#) | [Pax6](#) | [SHH](#) | [somatostatin](#) | [VIP](#)

FURTHER INFORMATION

Gord Fishell's homepage: <http://saturn.med.nyu.edu/research/dg/fishelllab/>
 Jean Hébert's homepage: http://neuroscience.aecom.yu.edu/faculty/primary_faculty_pages/hebert.html

ALL LINKS ARE ACTIVE IN THE ONLINE PDF